Foundations of gravitational waves in cosmology

Béatrice Bonga – 9 Dec 2021 – Synergies@Prague 2021 [BB+Prabhu, arXiv:2009.01243]

Radboud University



Invaluable tool: perturbation theory

 $g_{\mu\nu} = \overline{g}_{\mu\nu} + \epsilon h_{\mu\nu}$ Minkowski spacetime Kerr spacetime FLRW spacetime

But there is no canonical split!



From messy physics to peaceful realm



Different infinities



Key idea: bring infinity to a finite distance



Conformal diagram Minkowski



Asymptotic flatness



A physical spacetime $(\widehat{M}, \widehat{g}_{ab})$ is asymptotically flat if there exists a spacetime (M, g_{ab}) with boundary $\partial M \cong \mathcal{I} \cong \mathbb{R} \times \mathbb{S}^2$ such that

- 1. Ω and $g_{ab} = \Omega^2 \ \hat{g}_{ab}$ are smooth on M, $\Omega \ \widehat{=} \ 0$ and $n_a = \nabla_a \Omega$ is nowhere vanishing on \mathcal{I}
- 2. Einstein's equations are satisfied with \hat{T}_{ab} such that $\Omega^{-2}\hat{T}_{ab}$ has a smooth limit to \mathcal{I}

Consequences

- > Einstein's equation $\longrightarrow n^a$ is null on \mathcal{I}
- > q_{ab} = induced metric on \mathcal{I} is degenerate: 0 + +



This is common to <u>all</u> asymptotically flat spacetimes

$$\begin{cases} q_{ab}, n^{a} \dot{f} = \int w^{2} q_{ab}, w^{-1} n^{a} \dot{f} \end{cases}$$

Gravitational radiation is encoded in the next-order structure and differs from spacetime to spacetime

Nowhere in this construction did we introduce a split of the background and "gravitational waves".

The split occurs naturally at null infinity:

- universal structure is like a background,
- first order structure contains gravitational radiation,

and it is fully non-linear!



$$d\hat{s}^{2} = -UV \, du^{2} - 2U \, dudr + \gamma_{AB} (r \, d\theta^{A} + W^{A} \, du) (r \, d\theta^{B} + W^{B} \, du)$$

$$U = 1 + B/r^{2} + O(r^{-3}),$$

$$V = 1 - 2M/r + N/r^{2} + O(r^{-3}),$$

$$W^{A} = A^{A}/r + B^{A}/r^{2} + O(r^{-3}),$$

$$\gamma_{AB} = \Omega_{AB} + f_{AB}/r + \frac{1}{4}f^{2}\Omega_{AB}/r^{2} + O(r^{-3})$$

Generic asymptotically flat spacetime

Flat Space

Mass aspect

$$d\dot{s}^{2} = -UV \, du^{2} - 2U \, dudr + \gamma_{AB} (r \, d\theta^{A} + W^{A} \, du) (r \, d\theta^{B} + W^{B} \, du)$$

$$U = 1 + B/r^{2} + O(r^{-3}),$$

$$V = 1 - 2M/r + N/r^{2} + O(r^{-3}),$$

$$W^{A} = A^{A}/r + B^{A}/r^{2} + O(r^{-3}),$$

$$\gamma_{AB} = \Omega_{AB} + f_{AB}/r + \frac{1}{4}f^{2}\Omega_{AB}/r^{2} + O(r^{-3})$$

Radiative modes

Angular momentum aspect

Two definitions are equivalent!

Geometric description à la Penrose (with the conformal completion)

Coordinate description à la Bondi & Sachs

Asymptotic symmetry algebra

Spacetime diffeomorphism that leave the universal structure at null infinity invariant



Bondi-Metzner-Sachs algebra (BMS)

- Asymptotic symmetry algebra is *bigger* than Poincaré
- BMS = supertranslations & rotations

$$\begin{split} \xi^a \partial_a &= \begin{pmatrix} f(\theta,\varphi) + \frac{1}{2} & u \, D_A Y^A \end{pmatrix} \partial_u + Y^A \partial_A \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\$$

It provides quantities with a physical interpretation!



Critical assumption



Move far away from sources: 'spacetime becomes flat'

Expanding spacetimes are not asymptotically flat!



P. G. BERGMANN;

The only answer I can give is that the investigations date back less than two years, I believe, and that people have simply started with the mathematically simplest situation, or what they hoped was the simplest situation.

H. Bondi:

I regret it as much as you do, that we haven't yet got to the point of doing the Friedmann universe.

Conference Warsaw 1962

- Linear perturbation theory
 - Homogeneous solutions \rightarrow CMB
 - Inhomogeneous solutions from compact sources → geometric optics approximation

Goal:What can we learn from the full non-linear theory in this setting?

Expansion rates

Expanding spacetimes

Acceleration * Current universe * Λ > 0

 $\begin{array}{l} \textbf{Deceleration} \\ * \text{ Matter dominated era} \\ * \text{ Radiation dominated era} \\ * \Lambda = 0 \end{array}$

Radiation zones





 $\Lambda > 0$

Decelerating FLRW spacetimes

 $d\hat{s}^{2} = \alpha^{2}(\eta) \left[-d\eta^{2} + d\tau^{2} + r^{2} S_{AB} dx^{A} dx^{B} \right]$ $\begin{pmatrix} physical \\ physical$

$$S = \frac{2}{3(1+w)}$$
$$O \leq 5 < 1$$
$$-\frac{1}{3} < w < \infty$$

$$\frac{P = W P}{W = 1}$$

$$W = 1$$

$$W = 1/3$$

$$W = 0$$

$$W = 0$$

$$W = -1$$

$$W = -1$$

$$W = -1$$

$$d\hat{S}^2 = \alpha^2(\eta) \left[-d\eta^2 + dr^2 + r^2 S_{AB} dx^A dx^B \right]$$



Choose
$$\Omega = 2(\cos \frac{y}{2} \cos \frac{y}{2})^{\frac{1}{1-s}} (\sin \frac{U+V}{2})^{\frac{-s}{1-s}}$$

 $ds^2 = \Omega^2 ds^2 = -dUdV + \sin(\frac{V-U}{2})^2 S_{AB} dx^A dx^B$
 \longrightarrow Can add $V = -U = V = T_0$, because this
metric is smach everywhere including at the
boundaries



The conformal factor



Simple resolution

⊥ is smooth @J (" $* \ \ \, \underline{\mathcal{O}}^{l-2} \stackrel{<}{=} \bigcirc$ * $\nabla_{a} \Omega^{r-s} \neq O$

Define the normal to Y using 21-5 \implies $N_{a} = \frac{1}{1-S} \sqrt{a} S l^{-S}$ $= \Omega^{-S} \nabla_{A} \Omega$ $\hat{z} = \frac{2^{-S}}{1-S} \left(c \ll \frac{1}{2} \right)^{-S} \nabla_{a} V$

For asymptotically flat spacetimes, $\Omega^{-2}\hat{T}_{ab}$ should have a limit to \mathcal{I} but FLRW spacetimes are homogeneous, so there is matter *everywhere*!

$$\lim_{x \to 0} 8\pi G g^{ab} \widehat{T}_{ab} = \frac{68(1-s)}{(1-s)^2} \left(8\pi C \frac{U}{2}\right)^2 \rightarrow NON - VANISHING$$

$$8\pi G \widehat{T}_{ab} = 28 S 2^{2(s-1)} nah_b + 2S S^{s-1} T_{a} n_{b} + finite$$

$$universal$$

$$depends on doice S$$

$$T_{a} = ton \frac{U}{2} (T_{a}U + T_{a}V)$$

A physical spacetime $(\widehat{M}, \widehat{g}_{ab})$ admits a cosmological null asymptote if there exists a spacetime (M, g_{ab}) with boundary $\partial M \cong \mathcal{I} \cong \mathbb{R} \times \mathbb{S}^2$ such that

(1)
$$\Omega \cong 0$$
, Ω^{1-s} and $g_{ab} = \Omega^2 \hat{g}_{ab}$ is smooth on M ,
 $n_a = \Omega^{-s} \nabla_a \Omega$ is nowhere vanishing on \mathcal{I} (for $0 \le s < 1$)

(2) Einstein's equations are satisfied with
$$\hat{T}_{ab}$$
 such that

$$\lim_{\sigma \neq \mathscr{I}} g^{ab} \hat{T}_{ab} \quad \text{exists}$$

$$\lim_{\sigma \neq \mathscr{I}} \Omega^{1-s} \left[8\pi \hat{T}_{ab} - 2s \Omega^{2(s-1)} n_a n_b \right] \stackrel{\sim}{=} 2s \tau_{(a} n_{b)}$$

A physical spacetime $(\widehat{M}, \widehat{g}_{ab})$ admits a cosmological full asymptote if there exists a spacetime (M, g_{ab}) with boundary $\partial M \cong \mathcal{I} \cong \mathbb{R} \times \mathbb{S}^2$ such that

(1)
$$\Omega \cong 0, \Omega^{1-s}$$
 and $g_{ab} = \Omega^2 \hat{g}_{ab}$ is smooth on M ,
 $n_a = \Omega^{-s} \nabla_a \Omega$ is nowhere vanishing on \mathcal{I} (for $0 \le s < 1$)

(2) Einstein's equations are satisfied with \hat{T}_{ab} such that $\lim_{d \to \mathscr{I}} g^{ab} \hat{T}_{ab} \quad \text{exists}$ $\lim_{d \to \mathscr{I}} \Omega^{1-s} \left[8\pi \hat{T}_{ab} - 2s \Omega^{2(s-1)} n_a n_b \right] \stackrel{\frown}{=} 2s \tau_{(a} n_{b)}$

All smooth vector fields that map $2 q_{ab} = w^2 q_{ab}$, $n'^a = w^{-1-s} n^a f$

$$\implies b_s \cong \mathcal{BO}(1,3) \times \mathcal{S}_s$$

Didn't we know this already?

| es 🛛 Mendeley Desktop 👻 | Thu 11:10 Mendeley Deskton | ÷ 4≬ 8 ▼ |
|---------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| File Edit View Go Tools Help | | urch Béatrice 🚭 |
| Select Pan Note Highlight Color | Zoom Zoom To Fit Fullscreen Sync Help | |
| My Library 🖲 BMS in Cosmology 🗱 | | A |
| | BMS in Cosmology | |
| | A. Kehagias ^{a} and A. Riotto ^{b} | |
| | ^a Physics Division, National Technical University of Athens, 15780 Zografou Campus, Athens, Greece | : |
| 5 Oct 2016 | ^b Department of Theoretical Physics and Center for Astroparticle Physics (CAP) 24 quai E. Ansermet, CH-1211 Geneva 4, Switzerland | |
| [th] | Abstract | |
| ep-1 | Symmetries play an interesting role in cosmology. They are useful in characterizing the cosmo- | |

Page 1 of 33



Not exactly BMS in cosmology

| 🐱 Mendeley Desktop 🔻 | Thu 11:10 | ÷ 4ù 0 - | |
|------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|------------|
| le <u>E</u> dit <u>V</u> iew Go <u>Tools</u> <u>H</u> elp AT <u></u> Select Pan Note Highlight Color | Mendeley Desktop | Béatrice 😵 | |
| My Library 🖲 BMS in Cosmology | * | | |
| | s 🕱 Mendeley Desktop - Thu 15:53 | | ⊊ aù @ ▼ |
| | File File View Co. Table Viele | | 008 |
| | Line Joint grew on Jools grew All → (*) □ / (*) → (*) 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 | Q• Search | Béatrice 🎲 |
| | 🖾 My Library 📍 BMS in Cosmology 🗱 📍 BMS-like symmetries in co 🗱 | | |
| 2016 | BMS-like symmetries in cosmology Béatrice Bonga ^{1,*} and Kartik Prabhu [†] | Physics | |
| Oct | Radboud University, 6525 AJ Nijmegen, The Nether | rlands | - |
|] 5 | Abstract | | |
| -th | Null infinity in asymptotically flat spacetimes posses a rich mathematical s | structure; including the | |
| ep | BMS group and the Bondi news tensor that allow one to study gravitation | al radiation rigorously. | |
| | However, FLRW spacetimes are not asymptotically flat because their str | ess-energy tensor does | |
| | not decay sufficiently fast and in fact diverges at null infinity. This class radiation-dominated FLRW spacetimes. We define a class of spacetimes v infinity is similar to FLRW spacetimes: the stress-energy tensor is allow | s includes matter- and whose structure at null red to diverge and the | v |

notion of mass and linear momentum?

 $\Psi_4 := -C_{abcd}\overline{m}^a n^b \overline{m}^c n^d \stackrel{\circ}{=} 0, \quad \tilde{\Omega}^{-1} \Psi_4 \stackrel{\circ}{=} \left(\frac{1}{2} \partial_u^2 C_{AB} + s \partial_u \eth_A \tau_B + \frac{1}{2} s^2 \tau_A \partial_u \tau_B \right) \overline{m}^A \overline{m}^B$

$$\Psi_{3} := -C_{abcd}l^{a}n^{b}\overline{m}^{c}n^{d} \stackrel{\widehat{=}}{=} -\frac{s}{4}\partial_{u}\tau_{A}\overline{m}^{A}$$

$$\Psi_{2} := -C_{abcd}l^{a}m^{b}\overline{m}^{c}n^{d} \stackrel{\widehat{=}}{=} -\frac{1}{6}\left[W^{(2)} - 1 - s\left(\partial_{u}\tau + \frac{1}{2}\eth_{A}\tau^{A} + s\tau_{A}\tau^{A} + \frac{3}{2}i\epsilon^{AB}\eth_{A}\tau_{B}\right)\right]$$



Class of spacetimes at least as big as asymptotically flat spacetimes!



Linearization stability still open question!

Conclusion

- Gravitational radiation can be studied using asymptotics in the full non-linear theory
- Asymptotic symmetry algebra provides charges and fluxes with a physical interpretation
 - > Asymptotic flat spacetimes: BMS
 - Asymptotic cosmological null asymptotes: BMS-*like*, there is no translation subalgebra!





Others examples: Kerr-Newman spacetimes, Weyl spacetimes, etc.

How about FLRW?



No! This is similar to absence of a rotation subgroup for asymptotically flat spacetimes.